

Is The Enigmatic Source 3EG J1828+0142 a Galactic Microblazar ?

Y. Butt¹, presenting on behalf of: D. F. Torres^{2,3}, G. E. Romero^{4,5},
J. M. Paredes⁶, M. Ribó⁶, J. Martí⁷, J. Combi⁴, B. Punsly⁸

¹Harvard-Smithsonian Center for Astrophysics, Cambridge, MA, USA

²Department of Physics, Princeton University, Princeton, NJ, USA

³Lawrence Livermore National Laboratory, Livermore, CA, USA

⁴Instituto Argentino de Radioastronomía, Villa Elisa, B.A., Argentina

⁵presently at: CNRS/CEA/Saclay and Université de Paris, Paris, France

⁶Departament d'Astronomia i Meteorologia, Universitat de Barcelona, Spain

⁷Departamento de Física, Escuela Politécnica Superior, Universidad de Jaén, Spain

⁸4014 Emerald St., Torrance, CA, USA

Abstract :

The unidentified EGRET source 3EG J1828+0142, located at $(l, b) \approx (31.9, 5.8)$ (Hartman et al. 1999), is particularly enigmatic since it displays very high variability over timescales of months (Torres et al. 2000, Tompkins 1999) yet no blazar has been found within the 95% confidence contour (CC) of the gamma-ray location error-box (Punsly et al 2000). It also has an unusually steep γ -ray spectral index of $\Gamma = 2.76 \pm 0.39$. A multifrequency study of this source has been initiated and may reveal the presence of a novel type of compact galactic object: a microblazar. X-ray data, especially those of high spatial resolution from CHANDRA, would be particularly useful in establishing such a connection.

1 Introduction

The EGRET telescope onboard the Compton Gamma-Ray Observatory (CGRO) surveyed the entire sky in the 0.1 – 20 GeV γ -ray energy range and discovered 271 sources, of which roughly two-thirds still remain unidentified (Hartman et al. 1999). The source 3EG J1828+0142, located at $(l, b) \approx (31.9, 5.8)$ (Hartman et al. 1999), is particularly enigmatic since it displays very high variability over timescales of months (Torres et al. 2000, Tompkins 1999). However, no blazar has yet been found within the 95% confidence contour (CC) of the gamma-ray location error-box (Punsly et al 2000). 3EG J1828+0142's variability ranks 9σ above the average variability of known gamma-ray pulsars, making it the second most variable low-latitude non-transient γ -ray source. It also has an unusually steep γ -ray spectral index of $\Gamma = 2.76 \pm 0.39$. A multifrequency study of this source has been initiated and promises to reveal the presence of a novel type of compact galactic object.

2 Radio and X-rays

We have studied the region of the 3EG J1828+0142 error-box using radio data from the large-scale surveys by Haslam et al. (1981), Reich & Reich (1986) (see Combi et al. 2001) and small-scale VLA observations from the NVSS Sky Survey by Condon et al. (1998). In Fig. 1 we show a filtered map of the radio field at 1.4 GHz with the probability location contours of the 3EG source superposed. A large, shell-type structure can be clearly discerned. It is a weak radio source (the integrated flux density is 18.2 ± 2.1 Jy at 1.4 GHz, the non-thermal spectral index found for the radio emission is $\alpha = -0.72 \pm 0.18$), resembling a typical SNR. Its distance is ~ 940 pc making the radius of the remnant ~ 30 pc and its age $\sim 4 \times 10^4$ yr, assuming standard values for the particle density of the ISM and the original energy release (Punsly et al. 2000). Since the γ -ray source is located on the boundary of the remnant, the hypothetical compact source should have a birth velocity of ~ 700 km s⁻¹, in agreement with recent estimates (Lyne & Lorimer 1994). Fig. 1 shows the point-like radio sources de-

tected at 1.4 GHz with flux densities above 5 mJy. Most of them have no entry in any current catalog. We have computed spectral indices when possible, and upper limits for those sources not detected at 5 GHz in Condon et al.'s (1994) survey.

The region of interest has been observed (but with only ~ 340 sec exposure) in the ROSAT All-Sky Survey (RASS), and only a small corner (shown in the map) in pointed mode. We extracted all the ROSAT sources within 1 degree radius centered at $(l, b) = (31.8, 5.9)$ using the standard catalogs but none were found to be coincident with the radio sources in Fig. 1. An ASCA observation which includes part of the region of 3EG J1828+0142 is of too poor a spatial resolution to determine conclusively any possible radio-X-ray correspondences.

3 Modeling 3EG J1828+0142

The rapid variability of 3EG J1828+0142 clearly points to a compact object as the source of the γ -ray emission. But all identified EGRET AGNs are also strong radio sources (>1 Jy) with flat spectra, as expected from synchrotron jet-like sources where the γ -ray flux is the result of inverse Compton scattering. No source in the radio field seems to correspond to a typical γ -ray blazar. Independently, Mattox et al. 2001 also find it unlikely (from a statistical point of view) that this source could be an AGN. Although unexpected, discovering that the counterpart of the 3EG source is of extragalactic origin would be of great importance also: it could signify the existence of a new class of strong γ -ray emitting objects, which are radio quiet.

The galactic arena looks more promising, especially in light of the low Galactic latitude of the source. Recently, Kaufman-Bernado et al. (2002) have presented a model for γ -ray emitting microblazars (MB) based on external Compton scattering of thermal stellar photons from high-mass stellar companions. Vari-




Figure 1 : Upper panel: 1.4-GHz continuum map of the region around 3EG J1828+0142. Radio contours are shown in steps of 0.01 K starting at 0.04 K. The noise level is 0.02 K. Probability confidence contours (CC) for the location of the γ -ray source have been superposed in red. Lower panel: Point-like radio sources at the same frequency, detected by the VLA mostly within the 95% probability CC of the γ -ray source. The selected sources are within the 68% CC (Punsly et al. 2000). The position of the requested ACIS-S field of view is shown. The part of the circle in the lower-right corner was observed in pointed mode by ROSAT, and we discarded their possible association with the 3EG source due to poor spatial correlation. Sources #12, 15 and 16, and possibly also 8, are thermal. Sources #13, 19, 20 and 21 are outside the 68% CC, whereas #17 and 18 are outside even of the 95% contour.

ability is naturally introduced by the Newtonian precession of the accretion disk in the gravitational field of the star. This is expected to result in the precession of the relativistic jets, with the consequent flux variability due to variable Doppler boosting. Depending on the parameters of the model, such a MB can explain sources with a luminosity 10^{34-38} erg/s of γ -rays in the observer's frame. The non-thermal spectral energy distribution has two peaks, one from the synchrotron jet emission (at radio-IR energies) and other from Comptonization of stellar photons (at MeV-GeV energies). The total X-ray emission is $\sim 10^{34}$ erg s $^{-1}$. A model such as this may also explain the LS 5039/ 3EG J1824-1514 connection recently proposed by Paredes et al. (2000). In general, if the jets are not closely aligned to our line of sight, and particularly if the binary companion is not of high mass, the X-ray spectrum of a MQ will be affected by the disk emission. A power law plus an iron line was the case for LS5039 (Ribó et al. 1999). There are no known massive stars (spectral types B or O), superposed with any of the radio sources shown in Fig. 1. However, this absence could be explained by obscuration, since part of the region is in the line of sight to Serpens-Cauda Molecular Cloud Complex, lying at ~ 250 pc from Earth (Straizys et al. 1996).

Another possibility for explaining 3EG J1828+0142 is an isolated black hole (BH). It has been suggested that such objects could radiate γ -rays due to spherical accretion from the interstellar medium (Dermer 1997) or due to a charged magnetosphere (Punsly 1998, Punsly et al. 2000). The configuration of a simple axisymmetric magnetosphere around a maximally rotating BH attains a minimum energy configuration when the hole and the magnetosphere have equal and opposite charge. Punsly (1998) has shown that the magnetospheric charge can be supported in a stationary orbiting ring or disk. The entire magnetized system is stable only in an isolated environment, otherwise accretion onto the hole would disrupt the ring and its fields. Magnetized black holes (MBHs) are

charged similarly to neutron stars in pulsars, but unlike them, MBH have no solid surface and consequently no thermal X-ray emission is expected. These objects can support strong magnetized bipolar winds in the form of jets where γ -ray emission is originated by IC emission with synchrotron seed photons (Punsly 1998). Since both magnetic and rotation axes are always aligned in MBHs, their emission is non-pulsating (Punsly 1999). A detailed MBH model of the γ -ray emission from 3EG J1828+0142, consistent with its extreme variable behavior and steep spectral index, was presented in Punsly et al. (2000). The broad-band SED output of this model is shown in Fig. 2 of that paper. The expected luminosity of the MBH in the CHANDRA energy range is $\sim 10^{34}$ erg s $^{-1}$, with a spectral index $\Gamma = 1.5 \pm 0.25$. In the CHANDRA range, the spectrum should look like a featureless power law, there is no disk emission and no lines are expected. The X-ray source should appear as a point source positionally coincident with one of the non-thermal radio sources detected in the field, since the same leptonic population would also radiate synchrotron radio waves.

4 Future Observations

In order to locate any point-like X-ray emission coincident with the sources in the radio map we have requested a modest amount of CHANDRA time to explore the nature of sources 9, 11, and 14 in particular. These are the brightest non-thermal sources in the inner 68% CC of the gamma-ray source and are the most likely signatures of the putative compact galactic object. (This region was not observed in pointed mode by ROSAT.) CHANDRA's arcsecond spatial resolution will be extremely useful for this work since it will at once allow us to conclude the nature of the expected X-ray source (diffuse vs. point-like), and will yield a sufficiently precise location to permit the follow-up optical and VLBA studies which are being planned.

Combi J.A., Romero G.E., Benaglia P., Jonas J. 2001, A&A 366, 1047.

Condon J.J., et al., 1994, AJ 107, 1829.

Condon J.J., et al., 1998, AJ 115, 1693.

Dermer C.D., 1997, in Dermer C.D., Strickman M.S., Kurfess J.D. (eds.) Proceedings of the Fourth Compton Symposium, AIP, New York, p.1275.

Hartman R.C., et al., 1999, ApJS 123, 79.

Haslam C.G.T., et al., 1981, A&A 100, 209.

Kaufman Bernadó M.M., Romero G.E., Mirabel I.F., 2002, A&A, 385, L10.

Lyne A.G., Lorimer D.R., 1994, Nat 369, 127.

Mattox J. R., Hartman R. C., Reimer O., 2001, ApJS 135, 155.

Paredes J.M., Martí J., Ribó M., & Massi M. 2000, Sci, 288, 2340.

Punsly B., 1998, ApJ 498, 440.

Punsly B., 1999, ApJ 519, 336.

Punsly B., Romero G.E., Torres D.F., Combi J.A. 2000, A&A, 364, 552.

Reich W., Reich P., 1986, A&AS 63, 205.

Ribó M., Reig P., Martí J., Paredes J. M., A&A, 1999, 347, 518.

Romero G.E., Kaufman-Bernado M.M., Combi J., and Torres D.F., 2001 A&A 376, 599.

Straizys V., Cernis K., and Bartsiute S. 1996, Baltic Astronomy 5, 124,

Tompkins W., 1999, PhD Thesis, Stanford University, Stanford, [astro-ph/0202141].

Torres D.F., Romero G.E., Combi J.A., et al. 2001, A&A, 370, 468.

Torres D.F., Romero G.E., Eiroa E. 2002, ApJ, In press, [astro-ph/0112549].